

Hey'di AS
Bjørn Petersen
PB 13
2017 Frogner
Norway

03 December 2013
Report no. K130839-3
Page 1 of 2

CERTIFICATE OF ANALYSIS

Sample : Heydi Best Flyt

Methods : GEV version Feb. 23, 2011
ISO 16000-10, ISO 16000-3 and ISO 16000-6
Air exchange rate: 0.5h
Loading factor: 0.391 m²/m³

Result :

Substances	Result after 3 days	Result after 28 days
Methanal (formaldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
Ethanal (acetaldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
Ammoniak	< 0.2 µg/m ³	< 0.2 µg/m ³
Acetic acid	< 0.2 µg/m ³	< 0.2 µg/m ³
Ethanol	< 0.2 µg/m ³	< 0.2 µg/m ³
2-propanone (acetone)	< 0.2 µg/m ³	< 0.2 µg/m ³
Butanal (butyraldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
2-butenal (crotonaldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
3-methylbutanal (isovaleraldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
Propanal (propionaldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³
3-methylbenz-aldehyde	< 0.2 µg/m ³	< 0.2 µg/m ³
2-methylbenz-aldehyde	< 0.2 µg/m ³	< 0.2 µg/m ³
4-methylbenz-aldehyde	< 0.2 µg/m ³	< 0.2 µg/m ³
Pentanal (valeraldehyde)	< 0.2 µg/m ³	< 0.2 µg/m ³

Result :

Substances	Result after 3 days	Result after 28 days
VOC	< 0.5 µg/m ³	< 0.5 µg/m ³
*TVOC (C6 – C16)	33 µg/m ³	6 µg/m ³
*C16	33 µg/m ³	6 µg/m ³
TSVOC (C16 - C22)	< 0.5 µg/m ³	< 0.5 µg/m ³

No agents according to IARC group 1 (carcinogens) was found.

Analysed by:

Toxicon AB



Daniel Kuster, lab ing.

Approved by:

Toxicon AB



Ingrid Trulsson, QA